

Reference List

- AAP Committee on Fetus and Newborn. (2003). Apnea, Sudden Infant Death Syndrome, and Home Monitoring. *Pediatrics*, 111(4), 914-917.
- AAP Committee on Injury and Poison Prevention. (1999). Safe Transportation of Newborns at Hospital Discharge. *Pediatrics*, 104(4), 986-987.
- AAP Subcommittee on Hyperbilirubinemia. (2004). Management of Hyperbilirubinemia in the Newborn Infant 35 or More Weeks of Gestation. *Pediatrics*, 114(1), 297-316.
- AAP Task Force on Sudden Infant Death Syndrome. (2011). Technical report: SIDS and other sleep-related infant deaths: expansion of recommendations for a safe infant sleeping environment. *Pediatrics*, 128(5), e1341-e1367.
- Adamkin, D.H. and Committee on Fetus and Newborn. (2011). Clinical Report—Postnatal Glucose Homeostasis in Late-Preterm and Term Infants. *Pediatrics*, 127(5), 575-579.
- Association of Women's Health, Obstetric and Neonatal Nurses (AWHONN). (2010). Assessment & Care of the Late Preterm Infant Guideline, 2010. (<http://www.awhonn.org/awhonn/store/productDetail.do?productCode=LPI-2010>)
- Bhutani, V.K. and the Committee on Fetus and Newborn. (2011). Phototherapy to Prevent Severe Neonatal Hyperbilirubinemia in the Newborn Infant 35 or More Weeks of Gestation. *Pediatrics*, 128(4), e1046-e1052.
- Bull MJ and Engle WA. (2009). Safe transportation of preterm and low birth weight infants at hospital discharge. *Pediatrics*, 123(5), 1424-1429.
- Campbell, M.A. (2006) Development of a Clinical Pathway for Near-Term and Convalescing Premature Infants in a Level II Nursery. *Advances in Neonatal Care*, 6(3), 150-164.
- Engle, W.A., Tomashek, K.M., Wallman, C.; Committee on Fetus and Newborn, American Academy of Pediatrics. (2007). "Late-Preterm" Infants: A Population at Risk. *Pediatrics*, 120(6), 1390-1401.
- Harigopal, S., Satish, H.P., Taktak, A.F.G., Southern, K.W., Shaw, N.J. (2011). Oxygen Saturation Profile in Healthy Preterm Infants. *Archives of Disease in Childhood: Fetal and Neonatal Edition*, 96(5), F339-F342.
- Kemper, A.R., Mahle, W.T., Martin, G.R., Cooley, W.C., Kumar, P., Morrow, W.R., Kelm, K., Pearson, G.D., Glidewell, J., Grosse, S.D., Howell, R.R. (2011). Strategies for Implementing Screening for Critical Congenital Heart Disease. *Pediatrics*, 128(5), e1259-e1267.
- Kent, A.L., Meskell, S., Falk, M.C., Shadbolt, B. (2009). Normative blood pressure data in non-ventilated premature neonates from 28-36 weeks gestation. *Pediatric Nephrology*, 21(1), 141-146.
- Kerac, M., McGrath, M., Grijalva-Eternod, C., Bizouerne, C., Saxton, J., Bailey, H., Wilkinson, C., Hirsch, J., Blencowe, H., Shoham, J., Seal, A. (2010). Chapter 7: Review of breastfeeding assessment tools. In *Management of Acute Malnutrition in Infants (MAMI) Project. Technical Review: Current evidence, policies, practices & programme outcomes*. Accessed Jan 2, 2013.

<http://reliefweb.int/sites/reliefweb.int/files/resources/8A7E77D26B35660F492576F70010D7DF-mami-report-complete.pdf>

- Mahle, W.T., Newburger, J.W., Matherne, G.P., Smith, F.C., Hoke, T.R., Koppel, R., Gidding, S.S., Beekman, R.H., Grosse, S.D. (2009). Role of Pulse Oximetry in Examining Newborns for Congenital Heart Disease: A Scientific Statement from the AHA and AAP. *Pediatrics*, 124(2), 823-836.
- Morgan, J.A., Young, L., McCormick, F.M., McGuire, W. (2012). Promoting growth for preterm infants following hospital discharge. *Archives of Disease in Childhood: Fetal and Neonatal Edition*, 97(4):F295-F298.
- National Perinatal Association. (2012). Multidisciplinary Guidelines for the Care of Late Preterm Infants. (<http://www.nationalperinatal.org/lptguidelines/pdf/NPALatePretermGuidelines-11-12.pdf>)
- Oklahoma Infant Alliance. (2010). Caring for the Late Preterm Infant: A Clinical Practice Guideline. (http://www.oklahomainfantalliance.org/lpi_guidelines.html)
- Pejovic, B., Peco-Antic, A., Marinkovic-Eric, J. (2007). Blood pressure in non-critically ill preterm and full-term neonates. *Pediatric Nephrology*, 22(2), 249-257.
- Raju, T.N., Higgins, R.D., Stark, A.R., Leveno, K.J. (2006). Optimizing care and outcome for late-preterm (near-term) infants: A summary of the workshop sponsored by the National Institute of Child Health and Development. *Pediatrics*, 118(3), 1207-1214.
- Versmold, H.T., Kitterman, J.A., Phibbs, R.H., Gregory, G.A., Tooley, W.H. (1981). Aortic Blood Pressure During the First 12 Hours of Life in Infants with Birth Weight 610 to 4,220 Grams. *Pediatrics*, 67(5), 607-613.
- Wagner, C.L. and Greer, F.R. (2008). Prevention of Rickets and Vitamin D Deficiency in Infants, Children, and Adolescents. *Pediatrics*, 122(5), 1142-1152.
- Wisconsin Shine Project at <http://wisconsinshine.org/>
- Young, L., Morgan, J., McCormick, F.M., McGuire, W. (2012). Nutrient-enriched formula versus standard term formula for preterm infants following hospital discharge. *Cochrane Database of Systematic Reviews*, 3:CD004696.
- Zubrow, A.B., Hulman, S., Kushner, H., Falkner, B. (1995). Determinants of Blood Pressure in Infants Admitted to Neonatal Intensive Care Units: A Prospective Multicenter Study. *Journal of Perinatology*, 15(6), 470-479.

March 2013